

[Main Menu](#) [Home Page](#)

---

## Low Molecular Wt Heparins:

---

### - Discussion:

- anti-Xa to anti IIa ratio for enoxaparin is about 3 to 1 where as unfractionated heparin has a ratio of 1 to 1;
- hence LMWH has a greater inhibitory effect on factor Xa and has a lesser effect on thrombin;
- thrombocytopenia and platlet dysfunction seems to occur less often than w/ unfractionated herapin;

### - Adult Dose:

- 30 mg SQ bid;

### - Labs:

- because LMWH have minimal effect on thrombin, there is minimal elevation of the PTT;

### - Lovenox / Enoxaparin:

- subcutaneous enoxaparin 30 mg every 12 hours;
- onset of activity occurs after about 3 hours;
- it is advised to delay administration of Lovenox for 12-24 hours following surgery;
- renal excretion is the main route of removal, and therefore dose reduction is required w/ significant renal dysfu
- in a total hip arthroplasty study by Colwell et al (JBJS July 1999, Vol 81-A), the overall rate of throboembolic d lovenox versus 3.7 % for patients receiving adjusted dose coumadin;
- rate of clinically significant bleeding was 0.9%;
- lovenox was continued only for the length of the hospitalization which averaged about 7 days;
- in the study by Hans Klaus Breddin et al. 2001, the authors performed a multicenter, open-label study with blinded adjudication of end points w/ randomly assigned patients with acute DVT to one of three treatment regimens: intravenous administration of unfractionated heparin;
- subcutaneous administration of a low-molecular-weight heparin, reviparin, twice a day for one week; or subcutaneous administration of reviparin once a day for four weeks.
- primary end point was evidence of regression of the thrombus on venography on day 21; secondary end were recurrent venous thromboembolism, major bleeding within 90 days after enrollment, and death;
- of the patients receiving unfractionated heparin, 40.2 percent (129 of 321) had thrombus regression, as compared with 53.4 percent (175 of 328) of the patients receiving reviparin twice daily and 53.5 % (167 of 312) of the patients receiving reviparin once daily;
- w/ regard to thrombus regression, reviparin administered twice daily was significantly more effective than heparin (relative likelihood of thrombus regression, 1.28; 97.5 percent confidence interval, 1.08 to 1.52; reviparin administered once daily (relative likelihood, 1.29; 97.5 percent confidence interval, 1.08 to 1.52).
- in acute deep-vein thrombosis, reviparin regimens are more effective than unfractionated heparin in reduc
- reviparin is also more effective than unfractionated heparin for the prevention of recurrent thromboemboli
- ref: Effects of a Low-Molecular-Weight Heparin on Thrombus Regression and Recurrent Thromboembolism Hans Klaus Breddin et al. N Engl J Med 2001;344:626-31.

### - Pentasaccharides:

- in the report by G.G Alexander et al., the authors studied the effects of the pentasaccharide Org31540/SR901 a highly selective, indirect inhibitor of activated factor X, is the first of a new class of synthetic antithrombot
- in a double-blind study, patients were randomly assigned to postoperative administration of one of five da doses of Org31540/SR90107A, given once daily, or to 30 mg of enoxaparin, given every 12 hours;
- treatment was continued for 10 days or until bilateral venography was performed after a minimum of 5 da
- of 933 patients treated, 593 were eligible for the efficacy analysis;
- with Org31540/SR90107A a dose effect was observed (P=0.002), with rates of venous thromboembolism 6.7 %, 1.7 %, 4.4 %, and 0 % for the groups assigned to 0.75 mg, 1.5 mg, 3.0 mg, 6.0 mg, and 8.0 mg the drug, respectively, as compared with a rate of 9.4 % in the enoxaparin group;
- reduction in the risk of venous thromboembolism was 82 percent for the 3.0-mg Org31540/SR90107A grc 29 percent for the 1.5-mg group (P=0.51);
- enrollment in the 6.0-mg and 8.0-mg Org31540/SR90107A groups was discontinued because of bleeding
- major bleeding occurred 3.5 percent less frequently in the 0.75-mg group (P=0.01) and 3.0 percent less fr in the 1.5-mg group (P=0.05) than in the enoxaparin group (in which the rate was similar to that in the
- the authors concluded that Org31540/SR90107A, a synthetic pentasaccharide, has the potential to impro ratio for the prevention of venous thromboembolism, as compared with low-molecular-weight heparin.

- ref: A Synthetic Pentasaccharide for the Prevention of Deep-Vein Thrombosis after Total Hip Replacement  
G.G Alexander et al. The New England Journal of Medicine -- March 1, 2001 -- Vol. 344, No. 9

---

Low-molecular-weight heparinoid compared with warfarin for prophylaxis of DVT in patients who are operated on  
Prevention of DVT and PE after THR. Comparison of low-molecular-weight heparin and unfractionated heparin.  
Prevention of DVT after elective hip surgery. A randomized trial comparing low molecular weight heparin with star  
Dihydroergotamine/heparin in the prevention of deep-vein thrombosis after total hip replacement. A controlled, prc  
A randomized controlled trial of a low-molecular-weight heparin (enoxaparin) to prevent deep-vein thrombosis in p  
The thromboprophylactic effect of a low-molecular-weight heparin (Fragmin) in hip fracture surgery. A placebo-con  
Original Article: A Randomized Controlled Trial of a Low-Molecular-Weight Heparin (Enoxaparin) to Prevent Deep  
Comparison of Enoxaparin and Warfarin for the Prevention of Venous Thromboembolic Disease After Total Hip Ar  
CW Colwell et al JBJS July 1999, Vol 81-A, No 7.